

## 小鼠抗 RFC2 单克隆抗体

中文名称: 小鼠抗 RFC2 单克隆抗体

英文名称: Anti-RFC2 mouse monoclonal antibody

别 名: RFC40

储 存: 冷冻 (-20℃) 避光

宿 主: Mouse

抗 原: RFC2

相关类别: 一抗

反应种属: Human

标 记 物: Unconjugate

克隆类型: mouse monoclonal

### 技术规格

#### Background:

The elongation of primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins , proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also called activator 1, is a protein complex consisting of five distinct subunits of 145, 40, 38, 37, and 36.5 kD. This gene encodes the 40 kD subunit, which has been shown to be responsible for binding ATP. Deletion of this gene has been associated with Williams syndrome. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008].

<b>Applications:</b>	WB, FC
<b>Name of antibody:</b>	RFC2
<b>Immunogen:</b>	Fusion protein of human RFC2
<b>Full name:</b>	replication factor C (activator 1) 2, 40kDa (RFC2), transcript variant 2
<b>Synonyms:</b>	RFC40
<b>SwissProt:</b>	P35250
<b>WB Predicted band size:</b>	39 kDa
<b>WB Positive control:</b>	HepG2, Hela, SVT2, COS7, Jurkat, MDCK cell lysates
<b>WB Recommended dilution:</b>	200-1000